

The surface of a lattice polytope

Gábor Hegedüs

Johann Radon Institute for Computational and Applied Mathematics

February 26, 2010

Abstract

My main results are simple formulas for the surface area of d -dimensional lattice polytopes using Ehrhart theory.

1 Introduction

Throughout the paper a *lattice polytope* $P \subseteq \mathbb{R}^d$ is a polytope whose vertices have integral coordinates.

Let $S \subseteq \mathbb{R}^d$ be a subset of the Euclidean space $\mathbb{R}^d$. Let $G(S)$ denote the *lattice point enumerator* of the set S , the number of lattice (integral) points in S , i.e., $G(S) = |(S \cap \mathbb{Z}^d)|$.

Let $\mathcal{P}$ denote an arbitrary d -dimensional lattice polytope. In the following we denote by

$$\nu\mathcal{P} := \{n\underline{x} : \underline{x} \in P\}$$

the dilatate of $\mathcal{P}$ by the integer factor $\nu \geq 0$.

In 1962 E. Ehrhart proved (see e.g. [1, Chapter 3, Chapter 5], [6]) the following Theorem:

⁰ **Keywords.** Lattice polytopes, Ehrhart polynomial

2010 Mathematics Subject Classification. 52B20, 52C07, 11H06

Theorem 1.1 *Let $\mathcal{P}$ be a convex d -dimensional lattice polytope in the Euclidean space $\mathbb{R}^d$. Then there exists a unique polynomial (the Ehrhart polynomial)*

$$E_{\mathcal{P}}(x) := \sum_{i=0}^d e_i(\mathcal{P})x^i \in \mathbb{Q}[x],$$

which has the following properties:

(1) *For all integers $\nu \geq 0$,*

$$E_{\mathcal{P}}(\nu) = |(\nu\mathcal{P}) \cap \mathbb{Z}^d|.$$

(2) *The leading coefficient $e_d(\mathcal{P})$ of $E_{\mathcal{P}}(x)$ is $\text{vol}(\mathcal{P})$, the volume of $\mathcal{P}$.*

(3) *If $\text{int}(\mathcal{P})$ denotes the interior of $\mathcal{P}$, then the reciprocity law states that for all integers $\nu > 0$,*

$$E_{\mathcal{P}}(-\nu) = (-1)^d |(\nu \cdot \text{int}(\mathcal{P})) \cap \mathbb{Z}^d|. \quad (1)$$

(4) *The second leading coefficient $e_{d-1}(\mathcal{P})$ of $E_{\mathcal{P}}(x)$ is the half of the lattice surface area of $\mathcal{P}$:*

$$e_{d-1}(\mathcal{P}) = \frac{1}{2} \sum_{F \text{ facet of } \mathcal{P}} \frac{\text{vol}_{d-1}(F)}{\det(\text{aff } F \cap \mathbb{Z}^d)}.$$

Here $\text{vol}_{d-1}(\cdot)$ denotes the $(d-1)$ -dimensional volume and $\det(\text{aff } F \cap \mathbb{Z}^d)$ denotes the determinant of the $(d-1)$ -dimensional sublattice contained in the affine hull of F .

(5) *The constant coefficient $e_0(\mathcal{P})$ of $E_{\mathcal{P}}(x)$ is 1.*

Let $\mathcal{P}$ be a convex d -dimensional lattice polytope, which contains the origin of the lattice in its interior. We say that $\mathcal{P}$ is *reflexive* if the dual polytope $\mathcal{P}^*$ is a lattice polytope, where the *dual polytope* of $\mathcal{P}$ is defined as

$$\mathcal{P}^* := \{y \in \mathbb{R}^d : \langle x, y \rangle \geq -1 \text{ for all } x \in \mathcal{P}\}.$$

A. M. Kasprzyk proved in [9, Proposition 3.9.2] the following equivalent characterization of reflexive Fano polytopes:

Proposition 1.2 *Let $\mathcal{P}$ be a d -dimensional Fano polytope. Then $\mathcal{P}$ is reflexive iff*

$$\text{vol}(\mathcal{P}) = \frac{\text{surf } \mathcal{P}}{d}. \quad (2)$$

In 1899 G. A. Pick published his famous formula in [11]. Using this formula we can compute easily the area of a lattice polygon. Pick showed that the following expression gives the area of a simple lattice polygon Q :

$$\text{Area}(Q) = I + \frac{B}{2} - 1,$$

where B is the number of lattice points on the boundary of Q and I is the number of lattice points in the interior of Q .

This formula can be derived easily from Ehrhart Theorem 1.1 (see e.g. [4, Chapter 4]).

My main results are similar simple formulas for the surface area of 3-dimensional and 4-dimensional lattice polytopes using Ehrhart theory.

2 The main results

Let $\mathcal{P}$ be a convex d -dimensional lattice polytope in $\mathbb{R}^d$. Denote by $\text{surf}(\mathcal{P})$ the lattice surface area of $\mathcal{P}$:

$$\text{surf}(\mathcal{P}) := \sum_{F \text{ facet of } \mathcal{P}} \frac{\text{vol}_{d-1}(F)}{\det(\text{aff } F \cap \mathbb{Z}^d)}.$$

Let $i(\mathcal{P})$ and $b(\mathcal{P})$ denote the numbers $|\text{int}(\mathcal{P}) \cap \mathbb{Z}^d|$ and $|\partial(\mathcal{P}) \cap \mathbb{Z}^d|$, respectively.

Here $\partial(\mathcal{P})$ denotes the boundary of the polytope $\mathcal{P}$.

Theorem 2.1 *Let $\mathcal{P}$ be a convex d -dimensional lattice polytope in $\mathbb{R}^d$.*

Suppose that d is an odd number. Let $t := \frac{d-1}{2}$. Then define the matrix

$$\mathbf{A}(\mathcal{P}, d) := \begin{pmatrix} b(\mathcal{P}) - 2 & 1^{d-3} & \dots & 1^2 \\ b(2\mathcal{P}) - 2 & 2^{d-3} & \dots & 2^2 \\ \vdots & \vdots & \ddots & \vdots \\ b(t\mathcal{P}) - 2 & t^{d-3} & \vdots & t^2 \end{pmatrix}$$

and

$$\mathbf{D}(\mathcal{P}, d) := \begin{pmatrix} 1^{d-1} & 1^{d-3} & \dots & 1^2 \\ \vdots & \vdots & \ddots & \vdots \\ t^{d-1} & t^{d-3} & \dots & t^2 \end{pmatrix}$$

Then

$$\text{surf}(\mathcal{P}) = \frac{\det(\mathbf{A}(\mathcal{P}, d))}{\det(\mathbf{D}(\mathcal{P}, d))} \quad (3)$$

Suppose that d is an even number.

Let $t := \frac{d}{2}$. Then define the matrix

$$\mathbf{B}(\mathcal{P}, d) := \begin{pmatrix} b(\mathcal{P}) & 1^{d-3} & \dots & 1 \\ b(2\mathcal{P}) & 2^{d-3} & \dots & 2 \\ \vdots & \vdots & \ddots & \vdots \\ b(t\mathcal{P}) & t^{d-3} & \vdots & t \end{pmatrix}$$

and

$$\mathbf{D}(\mathcal{P}, d) := \begin{pmatrix} 1^{d-1} & 1^{d-3} & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ t^{d-1} & t^{d-3} & \dots & t \end{pmatrix}$$

Then

$$\text{surf}(\mathcal{P}) = \frac{\det(\mathbf{B}(\mathcal{P}, d))}{\det(\mathbf{D}(\mathcal{P}, d))} \quad (4)$$

Proof. Let

$$E_{\mathcal{P}}(x) := \sum_{i=0}^d e_i(\mathcal{P})x^i \in \mathbb{Q}[x],$$

denote the Ehrhart polynomial of the polytope $\mathcal{P}$.

First suppose that d is an odd number. Let $0 \leq k \leq \frac{d-1}{2}$. Then using Theorem 1.1

$$i(k\mathcal{P}) + b(k\mathcal{P}) = L_{\mathcal{P}}(k) = e_d(\mathcal{P})k^d + e_{d-1}(\mathcal{P})k^{d-1} + \dots + 1 \quad (5)$$

and

$$-i(k\mathcal{P}) = -L_{\mathcal{P}}(-k) = -e_d(\mathcal{P})k^d + e_{d-1}(\mathcal{P})k^{d-1} - \dots + 1 \quad (6)$$

Suming (5) and (6) we get that

$$b(k\mathcal{P}) = 2e_{d-1}(\mathcal{P})k^{d-1} + 2e_{d-3}(\mathcal{P})k^{d-3} + \dots + 2$$

i.e.,

$$b(k\mathcal{P}) - 2 = 2e_{d-1}(\mathcal{P})k^{d-1} + 2e_{d-3}(\mathcal{P})k^{d-3} + \dots + 2e_2(\mathcal{P})k^2$$

for each $0 \leq k \leq \frac{d-1}{2}$. Solving this linear equation system using Cramer's rule we get that

$$e_{d-1}(\mathcal{P}) = \frac{\det(\mathbf{A}(\mathcal{P}, d))}{2\det(\mathbf{D}(\mathcal{P}, d))}.$$

But using Theorem 1.1 (4) we get that

$$\text{surf}(\mathcal{P}) = \frac{e_{d-1}(\mathcal{P})}{2},$$

and we get our result for odd d .

Suppose that d is an even number. Let $t := \frac{d}{2}$. Let $0 \leq k \leq \frac{d}{2}$. Using Theorem 1.1

$$i(k\mathcal{P}) + b(k\mathcal{P}) = L_{\mathcal{P}}(k) = e_d(\mathcal{P})k^d + e_{d-1}(\mathcal{P})k^{d-1} + \dots + 1 \quad (7)$$

and

$$i(k\mathcal{P}) = L_{\mathcal{P}}(-k) = e_d(\mathcal{P})k^d - e_{d-1}(\mathcal{P})k^{d-1} + \dots + 1 \quad (8)$$

Subtracting (8) from (7) we get that

$$b(k\mathcal{P}) = 2e_{d-1}(\mathcal{P})k^{d-1} + 2e_{d-3}(\mathcal{P})k^{d-1} + \dots + 2e_1(\mathcal{P})k \quad (9)$$

for each $0 \leq k \leq \frac{d}{2}$. We can again solve this linear equation system using Cramer's rule, hence

$$e_{d-1}(\mathcal{P}) = \frac{\det(\mathbf{B}(\mathcal{P}, d))}{2\det(\mathbf{D}(\mathcal{P}, d))}. \quad (10)$$

Theorem 1.1 (4) implies that

$$\text{surf}(\mathcal{P}) = \frac{e_{d-1}(\mathcal{P})}{2}, \quad (11)$$

and we get our result from (10) and (11).

Examples.

If $d = 3$, then $\text{surf}(\mathcal{P}) = b(\mathcal{P}) - 2$.

In [9, Proposition 10.3.2] A. M. Kasprzyk proved this formula from Pick's Theorem.

If $d = 4$, then

$$\text{surf}(\mathcal{P}) = \frac{b(2\mathcal{P}) - 2b(\mathcal{P})}{6}.$$

If $d = 5$, then

$$\text{surf}(\mathcal{P}) = \frac{b(2\mathcal{P}) - 4b(\mathcal{P}) - 6}{12}$$

Remark. A. M. Kasprzyk called my attention for the following consequence of Theorem 2.1.

Corollary 2.2 *Let $\mathcal{P}$ be a convex d -dimensional Fano lattice polytope in $\mathbb{R}^d$. Suppose that d is an odd number. Let $t := \frac{d-1}{2}$. Then define the matrix*

$$\mathbf{A}(\mathcal{P}, d) := \begin{pmatrix} b(\mathcal{P}) - 2 & 1^{d-3} & \dots & 1^2 \\ b(2\mathcal{P}) - 2 & 2^{d-3} & \dots & 2^2 \\ \vdots & \vdots & \ddots & \vdots \\ b(t\mathcal{P}) - 2 & t^{d-3} & \vdots & t^2 \end{pmatrix}$$

and

$$\mathbf{D}(\mathcal{P}, d) := \begin{pmatrix} 1^{d-1} & 1^{d-3} & \dots & 1^2 \\ \vdots & \vdots & \ddots & \vdots \\ t^{d-1} & t^{d-3} & \dots & t^2 \end{pmatrix}$$

Then

$$\text{vol}(\mathcal{P}) = \frac{\det(\mathbf{A}(\mathcal{P}, d))}{d \cdot \det(\mathbf{D}(\mathcal{P}, d))} \quad (12)$$

Suppose that d is an even number.

Let $t := \frac{d}{2}$. Then define the matrix

$$\mathbf{B}(\mathcal{P}, d) := \begin{pmatrix} b(\mathcal{P}) & 1^{d-3} & \dots & 1 \\ b(2\mathcal{P}) & 2^{d-3} & \dots & 2 \\ \vdots & \vdots & \ddots & \vdots \\ b(t\mathcal{P}) & t^{d-3} & \vdots & t \end{pmatrix}$$

and

$$\mathbf{D}(\mathcal{P}, d) := \begin{pmatrix} 1^{d-1} & 1^{d-3} & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ t^{d-1} & t^{d-3} & \dots & t \end{pmatrix}$$

Then $\mathcal{P}$ is a reflexive polytope iff

$$\text{vol}(\mathcal{P}) = \frac{\det(\mathbf{B}(\mathcal{P}, d))}{d \cdot \det(\mathbf{D}(\mathcal{P}, d))}. \quad (13)$$

□

Proof.

Corollary 2.2 is the obvious consequence of Theorem 2.1 and Proposition 1.2. □

Acknowledgements. I am indebted to Josef Schicho and Alexander M. Kasprzyk for their useful remarks.

References

- [1] M. Beck, S. Robins, Computing the Continuous Discretely: Integer-Point Enumeration in Polyhedra, Springer, 2007
- [2] M. Beck, J. A. De Loera, M. Develin, J. Pfeifle, R.P. Stanley, Coefficients and roots of Ehrhart polynomials. *Integer points in polyhedra—geometry, number theory, algebra, optimization*, 15–36, *Contemp. Math.*, **374**, Amer. Math. Soc., Providence, RI, 2005.
- [3] H. S. M. Coxeter, Introduction to geometry. Reprint of the 1969 edition. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1989
- [4] D. A. Cox, Lectures on Toric Varieties, lecture notes available online on 'http://www3.amherst.edu/~dacox'
- [5] D. DeTemple, J. M. Robertson, The equivalence of Euler's and Pick's theorems. *Math. Teacher* **67** (1974), no. 3, 222–226.
- [6] E. Ehrhart, Sur les polyèdres rationnels homothétiques à n dimensions. *C. R. Acad. Sci. Paris* **254** (1962) 616–618
- [7] R. W. Gaskell, M. S. Klamkin, P. Watson, Triangulations and Pick's theorem. *Math. Mag.* **49** (1976), no. 1, 35–37.

- [8] C. Haase, I. V. Melnikov, The reflexive dimension of a lattice polytope. *Ann. Comb.* **10** (2006), no. 2, 211–217
- [9] A. M. Kasprzyk, Toric Fano Varieties and Convex Polytopes, PhD Thesis, 2006, available online on 'http://magma.maths.usyd.edu.au/users/kasprzyk/research/pdf/Thesis.pdf'
- [10] A. C. F. Liu, Lattice points and Pick's theorem. *Math. Mag.* **52** (1979), no. 4, 232–235.
- [11] G. Pick, Geometrisches zur Zahlenlehre, *Sitzungber Lotos* (Prague) **19** (1899), 311–319
- [12] Rosenholtz, Ira; Calculating Surface Areas from a Blueprint. *Math. Mag.* **52** (1979), no. 4, 252–256.
- [13] P. R. Scott, The fascination of the elementary. *Amer. Math. Monthly* **94** (1987), no. 8, 759–768